

DIGITAL DISEASE MAPPING OF CITRUS CANKER FROM SELECTED CITRUS ORCHARDS IN POTHOWAR

Amar Mehmood, Gulshan Irshad, Qamar-Uz-Zaman, M. Usman Raja, Noureen Zafar and Ms. Tehmina Akbar

Department of Plant Pathology. PMAS-Arid Agriculture University Rawalpindi, Pakistan.

Corresponding author; amarmeh188@gmail.com

ABSTRACT

Agriculture is one of the essential sectors for the survival of humankind. Adapting technology as well as digitalization is very crucial for the field of agriculture to benefit the farmer as well as the consumer. Crop growth and yield are essential aspects that influence the field of agriculture as well as farmer economically and socially. So, it is necessary to have close monitoring at various stages of crop growth to identify the diseases at right time. But, humans naked may not be sufficient and sometimes it would be misleading scenarios arise. In this aspect, automatic recognition and classification of various diseases of a specific crop are necessary for accurate identification. This thought gave inspiration for the present proposed framework

Pakistan confers with a broad range of agro-climatic position, diverse from tropical to temperate, allowing 20 different types of fruits to grow. Citrus is an important fruit within the economically important family *Rutaceae* and is cultivated in Pakistan on a 20.0461 thousand ha with an annual production of 2.29 million tons. According to intentional reports published during 2015, Pakistan stands 16th country of world for production of good quality of citrus crops. However, during 2016 CABI released a report indicating a serious disease of citrus fruits like canker,

causing higher losses in citrus production caused by gram negative bacteria *Xanthomonas axonopodis*. Conventionally method for plant diseases diagnosis using hand lenses till to isolation lab techniques are laborious and not predictive for fungicidal application time to time, and make some treatment to control the diseases. By applying Artificial Intelligence techniques 4 to 5 thousands picture of Citrus canker spots from different orchards from different plant parts fruit and leaves at different location from pothowar were obtained . Then resize the images and retain images in convolutional neural network (CNN) by using python as computer language were applied . A model of *Citrus Fruits Detection (Multi Classification).ipynb(CFD)* was developed, that is able to detects the Citrus canker disease on Citrus Plant, and provides data at its initial stages (Low Infection) or at final stage (Severe Infection).

Keywords: Citrus, *Xanthomonas axonopodis*, Artificial, Intelligence ,CNN

INTRODUCTION

The distribution of crop diseases can affect the economy badly. The manual diagnoses of crop diseases are time-consuming and risk errors. Digital revolution is reinventing agriculture, which integrates advanced technologies, digital tools, information, and communication technologies to enhance the opportunities for agriculture improvement and performance. Digital agriculture is currently emerging as a consequence of several technological developments in artificial intelligence, remote sensing, and robotic systems. Such systems allow farmers to provide broad, precise, and accessible traditional agricultural products at the national and regional levels, and boost yield and quality while limiting environmental impact. It can also provide ease to the farmers for detecting plant disease, pests and weeds (Adaskaveg and Foster, 2004 ; Aruoma et al., 2012).

Citrus is producing in all the four provinces of Pakistan , But mainly in Province Punjab five districts, Sargodha (Bhalwal), Toba Tek Singh, MandiBahauddin, Sahiwal, Khanewal, Cover total area under cultivation 60% and production of Citrus is 65%. (Deqiu, 1993; Devatkal, and Naveena, 2010).). In Pakistan area and production of fruit increased by 22.8% and 21%, respectively, in the year

2005 better growth received by the growers, 0.8 million hectares area under fruit trees. (MINFAL, 2006; Franke and Menz, 2006; Gerhards and Oebel, 2006). Among fruits, citrus was cultivated on 192,274 hectares with annual production of 2,458,381 tons during 2005. Since the best growth of Kinnow (*Citrus reticulata*.), Pakistan produced 1.62 million tones citrus (Jeliazkova and D. Percival, 2003), which increased to 2.4 million tons in 2014 and 2015. (FAO, 2016), its production has been fistedly increasing to fulfill the demand in the country and abroad (Khan, 1992; Gliever and Slaughter, 2001; González-Molina et al., 2010).

The citrus stand first in area and production among all fruits produced worldwide. It is highly prized and remunerative fruit cultivated in more than 50 countries worldwide (Kl, 1993; Lin et al., 2009; Luo et al., 2008). The major fruit producers are China, Brazil, USA, and Mexico and contribute 47% of total citrus world production (Youseif et al., 2014; Mukti and Biswas , 2019). Brazil is the biggest citrus fruit production country in the world and most of the oranges are produced here (Wali et al., 2013; Nutter et al., 2006; Parvat et al.,2017). Global production of citrus has been considerably increased during the past few years and reached 3.2 metric million tons during 2016-17. The Oranges accounts for about 50 million tons (USDA, 2017). The availability of Pakistani kinnow in the European market is irregular and is mainly used as a raw material for juice. The Pakistan is exporting only 10% of its total Kinnow production which can be increased thrice by providing local farming better knowledge about growth to post harvest stages . It exported 372,160 tons Kinnow during the year 2015-16 and major international markets were Afghanistan, Azerbaijan, Indonesia, Mauritius, Oman and Philippines (Trade Development Authority of Pakistan, 2017).

In scientific literature, several techniques have been proposed to tackle the complex challenges in agriculture, such as decision support systems, plant disease detection, and other artificial intelligence-based techniques. Deep learning has shown most promising results for agricultural image processing such as plant disease detection, pesticides detection, plant type classification, etc. For instance,

the study proposed the detection of *Fusarium* head blight disease in wheat crops (Sankaran, 2016) . Here, deep convolutional neural network (CNN) and image processing techniques are employed to detect the diseased part of wheat leaf images. The authors exploit Bayesian deep learning for approximating the probability density for crop disease detection problems. Another deep CNN-based work suggests deploying a pre-trained model learned from usual massive datasets, and transferring it into a specific task trained with their data, like VGG Net and Image Net (Selvaraj et al., 2006;).

An automated wheat disease diagnostic system implementable on mobile devices to conduct a real-time diagnosis is based on deep learning and multiple instances learning (MIL). Their method uses four deep learning models, VGG-FCN-VD16, VGG-FCN-S, VGG-CNN-S, VGG-CNN-VD16, and are implemented on the leaf images dataset. The accuracies of VGG-CNN-VD16 and VGG-CNN-S are 73.00%, and 93.27%, respectively. However, the suggested model cannot detect the last stage of disease of the plant. Authors of applied the neural network, support vector machine, and fuzzy classifier for plant disease detection problems. They suggested that there is a need to work on diseases stage identification and quantification, real-world applications, and the reliability of a fully automatic system in agricultural sector. proposes the detection of the *Fusarium* head blight, a wheat crop disease (Saville, 2006) . They developed a deep convolutional neural network (DCNN) capable of extracting distinct wheat stems from a single image with a complicated environment. They also suggest a new method for identifying *Fusarium* head blight infected regions in each spike. In training, the model accurately detects the crop's diseased part, and the mean average precision is 0.9201. The results are better than k-means and Otsu's methods. However, this model requires a large dataset to detect the diseased part more accurately (Shahzadi et al., 2016; Shaw and Royle, 1989) .

A novel plant leaf disease detection model bases on deep CNN is proposed in(Lin et al., 2009). Transfer learning and deep CNN are used for the leaf disease detection problem. The deep CNN model could accurately differentiate 38

different groups of diseased and healthy plants using leaf images with 96.46% accuracy (Sunny and Peter, 2016) and used texture-based segmentation and simpler linear iterative clustering (SLIC) to capture and recognize the diseases and pests at early stages in corn crops. Classification is done through binary support vector machine (BSVM) and multi-class support vector machine (MSVM). The accuracy achieved for pest detection is 52%, which can be extendable used ResNet-101, VGG-16, and ResNet-50 can see blurred images, and yolov3 for pest and disease detection in rice crop, and detected blurred boundaries and irregular shapes. However, the model showed poor performance for fewer features present in the image frames (Thangadurai and Padmavathi, 2015).

The results presented by Mukti and Biswas (2019) suggested the use of image acquisition, image preprocessing, image segmentation, feature extraction, and classification techniques for the ginger plant disease detection problem. The system is linked with a digital/web camera, allowing farmers to take images of plant leaves. The collected images are processed using image processing techniques to identify diseases symptoms, disease type and notify the farmers about disease type through global system for mobile communications (GSM) interface. Then, relay turns on the pump installed in the device to release medicine to the infected plant according to the infected disease. However, the study does not consider a standard dataset of the ginger plant leaf images, and some diseases of ginger plants and pests are not discussed (Zaman et al., 2011).

This study focuses on making an autonomous system that detects ginger plant diseases, pest patterns, and deficiency nutrients through deep artificial neural network and learning techniques, namely VGG-16, CNN and MobileNetV2 in real-time circumstances. The study also involves developing a large-scale ginger plant dataset based on different stages. We present the classification of various diseases and nutrient deficiencies, and investigate the pest patterns in the leaf images. In addition, we exhibit the performance and ability of the model to

predict diseases with high accuracy. This study hopes to present the first step towards deep learning-based citrus canker disease detection. This research study presents the following key contributions. • To develop a standard dataset of leaves and fruits of citrus at healthy stage, citrus canker at initial stage and citrus canker at final stage.

MATERIAL AND METHODS

Data Collection

Collection of the images of Citrus fruits and leaves were collected at different stages twice in a month from emergence of canker spots at Knoot Research Farm, Pir Mehr Ali Shah Arid Agriculture University Rawalpindi at Chakwal and Panj Katha village in Attock. The size of data was almost 4000 images of canker spots by using Sensor (Iphone8 mobile Camera with 12-megapixel wide-angle f/1.8 camera with a larger. Data was taken from a specific distance of 01ft from infected sample to image collecting devices.

Image Acquisition and Description

To detect canker spots in real-time, there were three stages: Healthy Citrus (Healthy images of Fruits and Leaves), Canker at Initial Level (Canker spots on leaves and fruits at Initial level), Canker at Final Stage (Canker Spots on leaves and fruits at Final Stage).

RESULTS AND DISCUSSION

After collecting data, images were processed as 1.5×1.4 -m subject of view PNG documents within the Google Colab Note Book in Google drive. A detailed description of dataset distribution is provided in Table 1.

Table 1. A detailed description of dataset distribution

| Sr. No | Stages of Citrus | Training 80% | | Total | Testing 20% |
|--------|------------------|--------------|-------------|-------|-------------|
| | | Leaves Data | Fruits Data | | |
| | | | | | |

| | | | | | |
|----|-------------------------|------|------|------|------|
| 01 | Citrus Healthy | 512 | 512 | 1024 | |
| 02 | Citrus Canker (Initial) | 512 | 512 | 1024 | 768 |
| 03 | Citrus Canker (Final) | 512 | 512 | 1024 | |
| 04 | Total | 1536 | 1536 | 3072 | 3840 |

Data Augmentation and Processing

Data augmentation is a method of creating new training data from field data. We apply domain-specific techniques to samples from the training data to generate unique and distinct training instances. In this study, we augment the images by rescaling, rotating the images, changing the width and height shifts, zooming the images, and doing the horizontal flip. All the images are renamed by python code, resized by the cv2 library, and converted into RGB images for further data processing. The dimensions of the images are $(150 \times 150 \times 3)$, height and width are 150 and 150, and 3 represents RGB channel (Red, Green, Blue).



Figure 1. Citrus at Healthy stage.



Figure2. Canker on citrus leaves



Figure 03:

Citrus Canker at Final Stage in the field.

Classification

This step trains images of Citrus Canker disease. We use 80% data for training and the remaining 20% data for testing a detailed description of the deep learning algorithms is provided in the following table 2.

Table 2. Detail description of images use during algorithm.

| Sr. No | Stages of Citrus | Training 80% | | Total | Testing 20% |
|---------------|-------------------------|---------------------|--------------------|--------------|--------------------|
| | | Leaves Data | Fruits Data | | |
| 01 | Citrus Healthy | 512 | 512 | 1024 | |
| 02 | Citrus Canker (Initial) | 512 | 512 | 1024 | 768 |
| 03 | Citrus Canker (Final) | 512 | 512 | 1024 | |
| 04 | Total | 1536 | 1536 | 3072 | 3840 |

CNN Model


CNN is widely employed in the field of study. Image is represented by a three dimensional matrix is presented to CNN. Then, the convolutional layer extracts the characteristics of the image. Convolutional layer also includes ReLU activation, which reduces all negative values to zero. After convolution, the pooling layer is utilized to minimize the spatial volume of the input image. Then max pooling is used to minimize the spatial volume of the input image, and the 4×4 dimension input has been reduced to 2×2 dimensions. Then there is a fully connected layer, and the last is the logistic layer. The label, which is one-hot encoded, is contained in the output layer. A sequential model is used with Relu activation function. Dropout rate is 0.2 to reduce the over-fitting of the algorithm and sigmoid is used in the last layer. The hyper parameters used for CNN are given in Table 3.

Table 3. Hyper parameters tuning used for CNN.

| | |
|------------------------|--|
| Dataset ratio | There are three classes for citrus canker disease detection, Citrus (Healthy Stage), Citrus canker (Initial Stage), Citrus canker (Final Stage) fruits and leaves. 80% data is used for training and 20% data for testing for all detection cases. |
| Pre-processing | Images are renamed and resized to 150×150 |
| Batch size | Batch size is 32 for training the CNN |
| Epochs | Epochs are set to 60 to train the CNN.algorithm |
| Learning rate | Learning rate of 0.001 is set for the proposed model |
| Optimization algorithm | The CNN model is trained by Adam optimizer. |

The convolutions are a fixed size of 150×150 RGB (Red Green Blue) images during training. The pre-processing performed here removes the typical RGB value computed on the training phase-out of each pixel. The image is processed using a stack of convolutional layers, which employ filters with a small field of 3×3 . It is more complex and has nonlinear effects but has fewer parameters. In one of the settings, 1×1 convolution filters are used, which may be thought of as a linear modification of the input channels. For 3×3 convolution layers, the convolution phase and spatial padding of convolution input are kept to 1 pixel, ensuring that the spatial resolution is retained after convolution.

Spatial pooling is helped by five max-pooling layers that follow part of the convolution layers. Max pooling is done with stride 2 across a 2×2 -pixel frame. After a stack of convolution layers, there are three fully connected (FC) layers. The first two each have 4096 channels, whereas the third uses 1000 way ILSVRC classification and hence has 1000 channels, one for each class. The Softmax layer is the last layer. However, because of binary classification study. The last layer used here is the sigmoid layer. In all networks, the configuration of the ultimately linked layers is the same.






Figure 4. Training Data of Citrus (Healthy), Citrus canker (Initial Stage), Citrus canker (Final Stage).

This section presents the results of pest pattern detection. The dataset was split into three classes' pest pattern and healthy having 3072 images of training and 768 images for testing. Accuracy for CNN is shown in Figure 5. The number of epochs was represented on the x-axis, and it can be defined as the algorithm will learn in the number of times of the entire dataset. The y-axis represents the accuracy of the models, and accuracy is the ratio of the number of accurate predictions to the overall number of correct predictions. VGG-16 achieves better validation accuracy of 92% in disease detection.




Figure 5. Accuracy of CNN when epoch size is 45 and training accuracy is 92%.

Algorithm CNN

shows the accuracy 92%, loss was 0.3%, precision was 95%.

CONCLUSION

Citrus plant plays an important role in agriculture and medical field. So, this research represents an approach for canker detection in citrus. Digital image processing techniques can be employed to detect canker infected leaf diseases with accuracy compared to the traditional methods. The proposed CNN based leaf disease identification model *Citrus Fruits Detection (Multi Classification).ipynb(CFD)* developed, that detects the Citrus canker disease on Citrus Plant, and also detects and gives data about it that is it at initial stage (Low Infection) or at final stage (Severe Infection). Mapping of citrus canker disease done from four orchards at two different locations in pothowar region. In future we can add the management factor at all the stages for the citrus canker. To early manage the disease and also develop an application. Also develop a Mobile Application for the access of each farmer or citrus grower, for early detection of citrus canker and its proper management.

REFERENCES

- Adaskave J. E and Förster H. (2014). Integrated postharvest strategies for management of Phytophthora brown rot of citrus in the United States. In *Post-harvest Pathology* . pp; 123-131). Springer, Cham.
- Aruoma O. I., Landes B., Ramful-Baboolall D., Bourdon E., Neergheen-Bhujun V., Wagner K. H and Bahorun T. (2012). Functional benefits of citrus fruits in the management of diabetes. *Preventive medicine*, 54; S12-S16.
- Deqiu F. (1993). Research progress on chemical assorting of mandarin orange (*Citrus reticulata*). *Hubei Agricultural Sciences (China)*.
- Devatkal S. K. and Naveena B. M. (2010). Effect of salt, kinnow and pomegranate fruit by-product powders on color and oxidative stability of raw ground goat meat during refrigerated storage. *Meat science*, 85(2); 306-311.

- FAO.(2016). Food and Agriculture Organization. Area and production of Citrus.
[http://www. faostat.fao.org](http://www.faostat.fao.org).
- Franke J. and Menz G. (2006, September).Detection of fungal infection in wheat with high-resolution multispectral data. In *Remote Sensing and Modeling of Ecosystems for Sustainability III* (6298), pp. 99-106.
- Gerhards R. and Oebel H. (2006). Practical experiences with a system for site-specific weed control in arable crops using real-time image analysis and GPS-controlled patch spraying. *Weed research*, 46(3); 185-193.
- Gliever C and Slaughter D. C. (2001).Crop versus weed recognition with artificial neural networks. *ASAE paper*, 01-3104.
- González-Molina E., Domínguez-Perles R., Moreno D. A. and García-Viguera C. (2010). Natural bioactive compounds of Citrus limon for food and health. *Journal of pharmaceutical and biomedical analysis*, 51(2); 327-345.
- Jeliazkova and D. Percival E. (2003).Effect of drought on ericoid mycorrhizae in wild blueberry (*Vacciniumangustifolium*Ait.). *Canadian journal of plant science*, 83(3); 583-586.
- .
- Khan I. A. (1992, December). Virus and virus like diseases of citrus. In *Proc. first International Seminar on Citriculture in Pakistan*, pp. 2-5.
- KL H. (1993).Use of geographical information systems in analyzing large area distribution and dispersal of rice insects in South Korea. *Korean journal of applied entomology*, 32(3); 307-316.
- Lin H. C., Hsu S. T. and Tzeng K. C. (2009). Histopathology and bacterial populations of atypical symptoms-inducing *Xanthomonasaxonopodispv. citri* strains in leaves of grapefruit and mexican lime. *Journal of Plant Pathology*, 18; 125-134.
- Liu L., Li Y., Li S., Hu N., He Y., Pong R. and Law M. (2012). Comparison of next-generation sequencing systems. *Journal of Biomed Biotechnology*, 2012; 251364.

- Luo W., Taylor M. C. and Parker S. R. (2008). A comparison of spatial interpolation methods to estimate continuous wind speed surfaces using irregularly distributed data from England and Wales. *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 28(7); 947-959.
- MINFAL. (2006). Agricultural statistics of Pakistan. Government of Pakistan, Ministry of Food, Agriculture and Livestock.
- Mukti I. Z. and Biswas D. (2019, December). Transfer learning based plant diseases detection using ResNet50. In *2019 4th International conference on electrical information and communication technology (EICT)* (pp. 1-6).IEEE.
- Nutter F. W., Esker P. D. and Netto R. A. C. (2006). Disease assessment concepts and the advancements made in improving the accuracy and precision of plant disease data. *European Journal of Plant Pathology*, 115(1); 95-103.
- Parvat A., Chavan J., Kadam S., Dev S. and Pathak V. (2017). A survey of deep-learning frameworks. In *2017 International Conference on Inventive Systems and Control (ICISC)* (pp. 1-7).IEEE.
- Sankaran S., Mishra A., Ehsani R. and Davis C. (2010). A review of advanced techniques for detecting plant diseases. *Computers and electronics in agriculture*, 72(1); 1-13.
- Saville S. P., Lazzell A. L., Bryant A. P., Fretzen A., Monreal A., Solberg E. O. and Milne G. T. (2006). Inhibition of filamentation can be used to treat disseminated candidiasis. *Antimicrobial Agents and Chemotherapy*, 50(10); 3312-3316.

- Selvaraj M. G., Vergara A., Ruiz H., Safari N., Elayabalan S., Ocimati W. and Blomme G. (2019). AI-powered banana diseases and pest detection. *Plant Methods*, 15(1); 1-11.
- Shahzadi R., Ferzund J., Tausif M. and Suryani M. A. (2016). Internet of things based expert system for smart agriculture. *International Journal of Advanced Computer Science and Applications*, 7(9).
- Shaw M. W. and Royle D. J. (1989). Estimation and validation of a function describing the rate at which *Mycosphaerella graminicola* causes yield loss in winter wheat. *Annals of applied Biology*, 115(3); 425-442.
- Sunny S. and Peter R. (2016). Detection of canker disease on citrus leaves using image processing. *International Journal of Computer Engineering and Applications*, 10(3), 129-134.
- Thangadurai K. and Padmavathi K. (2015). Citrus canker disease detection using genetic algorithm in citrus plants. *International Journal of trends in Research and development* 2(5); 434-437.
- USDA . (2017) C USDA - Citrus: World Markets and Trade, United States Department of Agriculture, Foreign Agricultural Service, available in.
- Wali S., Munir F. and Mahmood T. (2013). Phylogenetic studies of selected Citrus species based on chloroplast gene, rps14. *International Journal of Agriculture and Biology* 15(2), 357-361.
- Youseif S. H., El-Halwagi A., Sayed H. A. and El-Itriby H. A. (2014). Chemical analyses, antibacterial activity and genetic diversity assessment of some Egyptian Citrus spp. cultivars. *African Journal of Biotechnology*, 13(26).
- Zaman Q. U., Esau T. J., Schumann A. W., Percival D. C., Chang, Y. K., Read S. M. and Farooque A. A. (2011). Development of prototype automated variable rate sprayer for real-time spot-application of agrochemicals in wild blueberry fields. *Comptact Electronic Agriculture*, 76(2); 175-182.

Journal of Biomaterials and Bio products Technology (JBBT)

(An Open Assess International Journal) ISSN: 2790-2609 (Online)

ISSN: 2790-2595 (Print)



Review 's Report

Guidelines for reviewers

Please consider following guidelines

1. Reviewers are requested kindly do not disclose any information regarding manuscript under review
2. Assert all hidden conflict of interest, seeking advice from the Journal if they are not sure about anything constitute a relevant interest in the manuscript under question
3. Acknowledge that peer review is largely undertaken to carry out their fair share of reviewing and in a timely mode.

We sincerely thank you for timely evaluation of the attached manuscript

Manuscript Evaluation

| | |
|--------------|---|
| Email: | asma.ahmed@imbb.uol.edu.pk ; asma.ahmad.aridian@gamil.com |
| Title: | Dr. |
| First Name: | Asma |
| Last Name: | Ahmed |
| Affiliation: | Institute of Molecular Biology and Biotechnology (IMBB), The University of Lahore |
| Country: | Pakistan |

Reviewer's Information

Manuscript Information

| | |
|--------------------|---|
| Manuscript Number: | Vol.2,Iss3.September 2022 |
| Manuscript Title: | DIGITAL DISEASE MAPPING OF CITRUS CANKER FROM SELECTED CITRUS ORCHARDS IN POTHOWAR |

Evaluation Report

Kindly enter comments per section of the manuscript

| | |
|----------------------|--|
| 1.1 GENERAL COMMENT: | Abstract is according to article and good |
|----------------------|--|

| | |
|--------------------------|---------------------------------------|
| Introduction: | Introduction are satisfactory |
| Methodology: | According to methods explained |
| Results: | Well written |
| Discussion: | Good |
| Bibliography/References: | Ok |
| Others: | - |
| Decision: | Recommended for publication |

Recommendation

Kindly mark the appropriate column.

| | |
|---|------------|
| Accept As Is: | |
| Requires Minor Corrections: | yes |
| Requires Moderate Revision: | |
| Requires Major Revision: | |
| Reject on grounds of (Please be specific) | |




Signature of Reviewer

Journal of Biomaterials and Bio products Technology (JBBT)

(An Open Assess International Journal) ISSN: 2790-2609 (Online)

ISSN: 2790-2595 (Print)



Review's Report

Guidelines for reviewers

Please consider following guidelines

4. Reviewers are requested kindly do not disclose any information regarding manuscript under review
5. Assert all hidden conflict of interest, seeking advice from the Journal if they are not sure about anything constitute a relevant interest in the manuscript under question
6. Acknowledge that peer review is largely undertaken to carry out their fair share of reviewing and in a timely mode.

We sincerely thank you for timely evaluation of the attached manuscript

Manuscript Evaluation

| | |
|--------------|--|
| Email: | Farhanamalik277@gmail.com |
| Title: | Dr. Farhana Bibi/ Asst. Professor GAUS |
| First Name: | Dr. Farhana |
| Last Name: | Bibi |
| Affiliation: | Asst. Professor Grand Asian University Sialkot, Pakistan |
| Country: | Pakistan |

Reviewer's Information

| | |
|--------------------|---|
| Manuscript Number: | Volume 2, Issue 3, 2022. |
| Manuscript Title: | DIGITAL DISEASE MAPPING OF CITRUS CANKER FROM SELECTED CITRUS ORCHARDS IN POTHOWAI |

Manuscript Information

Evaluation Report

Kindly enter comments per section of the manuscript

| | |
|----------------------|---|
| 1.2 GENERAL COMMENT: | Overall this is a quite productive study that is showing development of cankers on citrus fruits, which could play an important role in developing agricultural measures to control the disease. However, manuscript needs minor revisions as follows: |
| Introduction: | Introduction is well represented |
| Methodology: | Grammar check required |

| | |
|--------------------------|---|
| Results: | Typographic errors should be checked. Significant re be high lighted |
| Discussion: | Discussion needs to be more elaborated with recent st published |
| Bibliography/References: | Addition of more recent references required |
| Others: | |
| Decision: | Manuscript is acceptable after minor revisions |

Recommendation

| | |
|---|---|
| Accept As Is: | |
| Requires Minor Corrections: | √ |
| Requires Moderate Revision: | |
| Requires Major Revision: | |
| Reject on grounds of (Please be specific) | |

Kindly mark the appropriate column.



Signature of Reviewer

paper

by Maham misbah

Submission date: 02-Nov-2022 12:26AM (UTC+1100)

Submission ID: 1941388346

File name: duplicate.docx (24.71K)

Word count: 2655

Character count: 14349

DIGITAL DISEASE MAPPING OF CITRUS CANKER FROM SELECTED CITRUS ORCHARDS IN POTHOWAR

ABSTRACT

For humanity to survive, agriculture is among the most important industries. For the area of agriculture to serve both the farmer and the consumer, technology and digitalization must be used. Crop production and growth are important factors that have an impact on the agricultural industry as well as the farmer's social and economic well-being. In order to detect illnesses at the appropriate time, careful monitoring at different phases of crop development is required. But human beings in their natural state may not be adequate, and false scenarios might sometimes appear. For precise identification in this regard, automated detection and categorization of many illnesses affecting a particular crop are required. This idea served as the motivation for the current framework proposal.

Pakistan confers with a broad range of agro-climatic position, diverse from tropical to temperate, allowing 20 different types of fruits to grow. Citrus is an important fruit within the economically important family *Rutaceae* and is cultivated in Pakistan on a 20.0461 thousand ha with an annual production of 2.29 million tons. According to intentional reports published during 2015, Pakistan stands 16th country of world for production of good quality of citrus crops. However, during 2016 CABI released a report indicating a serious disease of citrus fruits like canker, causing higher losses in citrus production caused by gram negative bacteria *Xanthomonas axonopodis*. Conventionally method for plant diseases diagnosis using hand lenses till to isolation lab techniques are laborious and not predictive for fungicidal application time to time, and make some treatment to control the diseases. By applying Artificial Intelligence techniques 4 to 5 thousands picture of Citrus canker spots from different orchards from different plant parts fruit and leaves at different location from pothowar were obtained . Then resize the images and retain images in convolutional neural network (CNN) by using python as computer language were applied . A model of *Citrus Fruits Detection (Multi Classification).ipynb(CFD)* was developed, that is able to detects the Citrus canker disease on Citrus Plant, and provides data at its initial stages (Low Infection) or at final stage (Severe Infection).

Keywords: Citrus, *Xanthomonas axonopodis*, Artificial, Intelligence ,CNN

INTRODUCTION

Spreading agricultural diseases may have a negative impact on the economy. Manual crop disease diagnosis is labor-intensive and prone to mistakes. Technology advancements, digital

tools, information and communication technology, and the rise of the Internet have all contributed to a new era of agriculture that promises greater efficiency and productivity. As a result of recent breakthroughs in areas like as AI, remote sensing, and robotic technology, a new sector of the agricultural industry known as "digital agriculture" is developing. Farmers may increase productivity and quality while reducing environmental effect by using such systems, making traditionally grown goods available on a national and regional scale. In addition, it may make it simpler for farmers to identify plant diseases, pests, and weeds (Adaskaveg and Foster, 2004 ; Aruoma et al., 2012).

Citrus is producing in all the four provinces of Pakistan , But mainly in Province Punjab five districts, Sargodha (Bhalwal), Toba Tek Singh, MandiBahauddin, Sahiwal, Khanewal, Cover total area under cultivation 60% and production of Citrus is 65%. (Deqiu, 1993; Devatkal, and Naveena, 2010).). In Pakistan area and production of fruit increased by 22.8% and 21%, respectively, in the year 2005 better growth received by the growers, 0.8 million hectares area under fruit trees. (MINFAL, 2006; Franke and Menz, 2006; Gerhards and Oebel, 2006). Among fruits, citrus was cultivated on 192,274 hectares with annual production of 2,458,381 tons during 2005. Since the best growth of Kinnow (*Citrus reticulata*.), Pakistan produced 1.62 million tones citrus (Jeliaskova and D. Percival, 2003), which increased to 2.4 million tons in 2014 and 2015. (FAO, 2016), its production has been fistedly increasing to fulfill the demand in the country and abroad (Khan, 1992; Gliever and Slaughter, 2001; González-Molina et al., 2010).

On a global scale, citrus dominates in terms of both land used and yield. It's a lucrative crop that's grown in more than 50 nations across the globe because of how desirable and lucrative the fruit is (KI, 1993; Lin et al., 2009; Luo et al., 2008). China, Brazil, the United States, and Mexico are the world's four largest citrus fruit producers, accounting for 47% of worldwide output (Youseif et al., 2014; Mukti and Biswas , 2019). The majority of the world's oranges are harvested in Brazil, which is the largest producer of citrus fruits (Wali et al., 2013; Nutter et al., 2006; Parvat et al., 2017). The citrus crop has shown significant growth in recent years, with global output reaching 3.2 million tonnes in 2016–17. About 50 million metric tonnes of oranges are produced annually (USDA, 2017). Pakistani kinnow has sporadic and mostly unrefined availability on the European market and is primarily utilised as a raw ingredient for juice. By teaching local farmers more about the crop's development and post-harvest care, Pakistan might triple the amount of kinnow exported

as now. As of the end of the 2015–16 fiscal year, the country has sent out a total of 372,160 metric tons of Kinnow to various overseas markets (Trade Development Authority of Pakistan, 2017).

Several methods, such as decision support systems, plant disease detection, and other AI-based methods, have been presented in the scientific literature to address the complex problems in agriculture. In agriculture, deep learning has shown the greatest promise for image processing tasks including identifying plant diseases and herbicides and classifying plant types. For instance, the research recommended that wheat harvests be monitored for Fusarium head rot (Sankaran, 2016). Here, we use a deep convolutional neural network (CNN) and other image processing methods to identify the unhealthy areas in photographs of wheat leaves. Bayesian deep learning is used by the authors of to approximate the probability density for agricultural disease detection issues. Another piece of work based on deep CNNs proposes using a model that has already been trained on the typical large datasets and applying it to a new job using that dataset's data, such as VGG Net or Image Net (Selvaraj et al., 2006;).

With the use of deep learning and multi-instance learning, we can develop a smartphone app for diagnosing wheat diseases in real time (MIL). Four different deep learning models (VGG-FCN-VD16, VGG-FCN-S, VGG-CNN-S, VGG-CNN-VD16) are used in their approach, and the models are tested on the leaf pictures dataset. Accuracy-wise, VGG-CNN-VD16 is at 73,000%, whereas VGG-CNN-S is at 93,27%. However, the latter stages of plant disease are undetectable by the proposed approach. In order to solve plant disease identification issues, the authors of this study used a neural network, a support vector machine, and a fuzzy classifier. They advocated for more research into the detection and quantification of disease stages, practical applications, and the dependability of an automated system in agriculture. offers a method for identifying the wheat crop disease Fusariumhead blight (Saville, 2006). They used a deep convolutional neural network (DCNN) to separate individual wheat stems from their complex backgrounds in a single picture. They also propose a novel strategy for pinpointing the diseased areas of each spike caused by Fusarium head blight. The mean average accuracy of the model in identifying the sick portion of the crop during training is 0.9201. Both k-means and Otsu's approaches are outperformed by these findings. This approach, however, needs a large dataset in order to reliably identify the afflicted tissue (Shahzadi et al., 2016; Shaw and Royle, 1989).

A new model for detecting diseases in plant leaves, based on deep CNN, is suggested in (Lin et al., 2009). The challenge of identifying leaf diseases is addressed by using transfer learning and

deep CNN. Using texture-based segmentation and simpler linear iterative clustering (SLIC), the deep CNN model correctly distinguished between 38 groups of infected and healthy plants using leaf pictures (Sunny and Peter, 2016). This allowed for early detection of illnesses and pests in corn harvests. Two popular methods for doing classification are the binary support vector machine (BSVM) and the multi-class support vector machine (MSVM). The success rate of pest detection is 52%, which is a number that may be improved upon. Pest and disease identification in rice crops using ResNet-101, VGG-16, and ResNet-50 (which can see blurred pictures) and yolov3 (which can recognise irregular forms) (Li, D. et al., 2020). However, the model's performance was subpar when there were fewer characteristics in the images (Thangadurai and Padmavathi, 2015).

Mukti and Biswas (2019) reported research findings that supported the use of image capture, image preprocessing, picture segmentation, feature extraction, and classification methods to the disease detection issue in ginger plants. As the device is connected to a digital or web camera, growers may snap pictures of their plants' leaves. Images are analysed by image processing software to determine the nature of the illness, its symptoms, and how to best warn farmers through the global system for mobile communications (GSM) interface. Once the diseased plant's illness has been identified, the device's relay activates the pump housing the necessary medication. A standardised collection of ginger plant leaf photos was not used in the research, and neither were any of the known illnesses nor pests that affect ginger plants (Zaman et al., 2011).

The goal of this research was to develop an autonomous system that employs deep artificial neural networks and learning algorithms, such as VGG-16, CNN, and MobileNetV2, to identify illnesses, insect patterns, and nutritional deficiencies in ginger plants in real time. Creating a comprehensive dataset of ginger plants at various developmental phases is also part of the research objective. We provide a system for categorising illnesses and nutritional deficits and conduct research into pest patterns seen in digital photographs of leaves. Furthermore, we demonstrate the model's capabilities and effectiveness in accurately predicting illnesses. This research aspires to provide the first step toward the use of deep learning for the identification of citrus canker disease. The following are the main advantages of this study. In order to create a uniform data collection including images of citrus plants at three different stages of development (healthy, early canker, and late canker).

MATERIAL AND METHODS

Data Collection

Collection of the images of Citrus fruits and leaves were collected at different stages twice in a month from emergence of canker spots at Knoot Research Farm, Pir Mehr Ali Shah Arid Agriculture University Rawalpindi at Chakwal and Panj Katha village in Attock. The size of data was almost 4000 images of canker spots by using Sensor (Iphone8 mobile Camera with 12-megapixel wide-angle f/1.8 camera with a larger. Data was taken from a specific distance of 01ft from infected sample to image collecting devices.

Image Acquisition and Description

To detect canker spots in real-time, there were three stages: Healthy Citrus (Healthy images of Fruits and Leaves), Canker at Initial Level (Canker spots on leaves and fruits at Initial level), Canker at Final Stage (Canker Spots on leaves and fruits at Final Stage).

RESULTS AND DISCUSSION

After collecting data, images were processed as 1.5×1.4 -m subject of view PNG documents within the Google Colab Note Book in Google drive. A detailed description of dataset distribution is provided in Table 1.

Data Augmentation and Processing

Data augmentation is a way to generate more training data from existing field data. To produce novel and diverse training examples, we use domain-specific methods on subsets of the training data. In this research, we modify the photographs in many ways, including resizing, rotating, shifting horizontally and vertically, zooming in, and inverting. The photos are reorganized for further processing by having their filenames changed using python code, having their sizes adjusted by the cv2 library, and having their color spaces transformed to RGB. Image height and width are both 150 pixels, while the third digit, 3, indicates the RGB channel size (Red, Green, Blue).

Classification

This process educates mental pictures of Citrus Canker. The following table 2 provides a full overview of the deep learning techniques that we use, and we utilize 80% of the available data for training and 20% for testing.

CNN Model

Research institutions often use CNN. The image is delivered to the CNN as a 33 matrix. This information is then sent to the convolutional layer, which is responsible for extracting the image's unique properties. All negative values are normalized to zero thanks to the ReLU activation that is part of the convolutional layer. The pooling layer is used to reduce the input image's spatial volume after convolution. After that, we apply max pooling to decrease the input image's spatial volume from 4 by 4 to 2 by 2, thereby eliminating the fourth dimension. A completely linked layer follows, and finally the logistics layer. In the output layer, you'll find the label, which has been encoded using one-hot encoding. The Relu activation function is used to a sequential model. A 0.2 percentage point dropout rate and a sigmoid activation function in the last layer help prevent the algorithm from being over-fit. Table 3 lists the many CNN hyper parameters that have been tested.

During the training process, the convolutions have a fixed size of 150 x 150 RGB (Red Green Blue) pictures. The usual RGB value generated on the training phase-out of each pixel is discarded during this preprocessing. Images are processed by a series of convolutional layers, each of which uses filters with a 3 x 3 pixel field size. Complexity and nonlinear effects are increased while the number of parameters is reduced. One option makes use of 11 convolution filters, which may be seen as a linear transformation of the input channels. The spatial resolution is retained after the convolution for 3x3 convolution layers as the convolution phase and spatial padding of convolution input are preserved to 1 pixel.

After some of the convolution layers, there are five max-pooling layers that aid with spatial pooling. For a 2 by 2 image, the maximum pooling stride is 2. Following the series of convolution layers are three fully connected (FC) layers. For example, each of the first two have 4096 channels, but the third employs 1000-way ILSVRC classification and hence needs 1000 channels, one for each class. The very final layer is a Softmax layer. In spite of this, research using binary categories has limitations. After all other layers have been used, the sigmoid layer is employed. All networks have the same structure of interconnected layers.

The outcomes of pest pattern identification are shown here. 3072 photos were used for training and 768 were used for testing, with the dataset being divided into three classes representing the pest pattern and the healthy state. Figure 5 displays the precision of CNN. The x-axis showed the number of epochs, which can be thought of as the number of times the algorithm would learn from the complete dataset. The y-axis shows how well each model can make predictions, with accuracy defined as the proportion of right forecasts that are really correct. In the case of illness diagnosis, VGG-16 has a higher validation accuracy of 92%.

CONCLUSION

Citrus plant plays an important role in agriculture and medical field. So, this research represents an approach for canker detection in citrus. Digital image processing techniques can be employed to detect canker infected leaf diseases with accuracy compared to the traditional methods. The proposed CNN based leaf disease identification model *Citrus Fruits Detection*

(Multi Classification).ipynb(CFD) developed, that detects the Citrus canker disease on Citrus Plant, and also detects and gives data about it that is it at initial stage (Low Infection) or at final stage (Severe Infection). Mapping of citrus canker disease done from four orchards at two different locations in pothowar region. In future we can add the management factor at all the stages for the citrus canker. To early manage the disease and also develop an application. Also develop a Mobile Application for the access of each farmer or citrus grower, for early detection of citrus canker and its proper management.

paper

ORIGINALITY REPORT

10%

SIMILARITY INDEX

3%

INTERNET SOURCES

10%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

| | | |
|---|--|-----|
| 1 | Hamna Waheed, Noureen Zafar, Waseem Akram, Awais Manzoor, Abdullah Gani, Saif ul Islam. "Deep Learning Based Disease, Pest Pattern and Nutritional Deficiency Detection System for "Zingiberaceae" Crop", Agriculture, 2022 Publication | 6% |
| 2 | Submitted to Babes-Bolyai University Student Paper | 1% |
| 3 | Babak Maser, Andreas Uhl. "Using CNNs to Identify the Origin of Finger Vein Sample Images", 2021 IEEE International Workshop on Biometrics and Forensics (IWBF), 2021 Publication | 1% |
| 4 | usave.co.uk Internet Source | <1% |
| 5 | Submitted to University of East London Student Paper | <1% |
| 6 | researcherslinks.com Internet Source | <1% |

7

Ram Krishn Mishra, Siddhaling Urolagin, J. Angel Arul Jothi, Pramod Gaur. "Deep hybrid learning for facial expression binary classifications and predictions", Image and Vision Computing, 2022

Publication

<1 %

8

Takaaki Fujiwara, Ayumi Fujishima, Yui Nakamura, Kenji Tajima, Min Yao. "Structural snapshot of a glycoside hydrolase family 8 endo- β -1,4-glucanase capturing the state after cleavage of the scissile bond", Acta Crystallographica Section D Structural Biology, 2022

Publication

<1 %

9

link.springer.com

Internet Source

<1 %

10

Vijaypal Singh Dhaka, Sangeeta Vaibhav Meena, Geeta Rani, Deepak Sinwar, Kavita, Muhammad Fazal Ijaz, Marcin Woźniak. "A Survey of Deep Convolutional Neural Networks Applied for Prediction of Plant Leaf Diseases", Sensors, 2021

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On